

Science News

from research organizations



Probiotic use is a link between brain foginess, severe bloating

Date: August 6, 2018

Source: Medical College of Georgia at Augusta University

Summary: Probiotic use can result in a significant accumulation of bacteria in the small intestine that can result in disorienting brain foginess as well as rapid, significant belly bloating, investigators report.

Share: [f](#) [t](#) [p](#) [in](#) [✉](#)

FULL STORY

Probiotic use can result in a significant accumulation of bacteria in the small intestine that can result in disorienting brain foginess as well as rapid, significant belly bloating, investigators report.

In a published study of 30 patients, the 22 who reported problems like confusion and difficulty concentrating, in addition to their gas and bloating, were all taking probiotics, some several varieties.

When investigators looked further, they found large colonies of bacteria breeding in the patients' small intestines, and high levels of D-lactic acid being produced by the bacteria lactobacillus' fermentation of sugars in their food, says Dr. Satish S.C. Rao, director of neurogastroenterology/motility and the Digestive Health Clinical Research Center at the Medical College of Georgia at Augusta University.


D-lactic acid is known to be temporarily toxic to brain cells, interfering with cognition, thinking and sense of time. They found some patients had two to three times the normal amount of D-lactic acid in their blood. Some said their brain foginess -- which lasted from a half hour to many hours after eating -- was so severe that they had to quit their jobs.

The report in the journal *Clinical and Translational Gastroenterology* appears

Most Popular

this week

HEALTH & MEDICINE



'AeroNabs' Promise Powerful, Inhalable Protection Against COVID-19

COVID-19 False Negative Test Results If Used Too Early

The Six Strains of SARS-CoV-2

MIND & BRAIN

New Study Suggests ADHD Like Behavior Helps Spur Entrepreneurial Activity

Loss of Smell and Taste Validated as COVID-19 Symptoms in Patients With High Recovery Rate



'Little Brain' or Cerebellum Not So Little After All

LIVING & WELL

Boy or Girl? It's in the Father's Genes

Study Predicts Millions of Unsellable Homes Could Upend Market

The Best Material for Homemade Face Masks May Be a Combination of Two Fabrics

Strange & Offbeat

HEALTH & MEDICINE

Swallowing This Colonoscopy-Like Bacteria Grabber Could Reveal Secrets About Your Health

Evidence in Mice That

to be the first time the connection has been made between brain foginess, bacterial overgrowth in the small intestine, high levels of D-lactic acid in the gut and probiotic use, Rao says.

"What we now know is that probiotic bacteria have the unique capacity to break down sugar and produce D-lactic acid. So if you inadvertently colonize your small bowel with probiotic bacteria, then you have set the stage for potentially developing lactic acidosis and brain foginess," Rao says.

While probiotics can be beneficial in some scenarios, like helping a patient restore his gut bacteria after taking antibiotics, the investigators advised caution against its excessive and indiscriminate use.

"Probiotics should be treated as a drug, not as a food supplement," Rao says, noting that many individuals self-prescribe the live bacteria, which are considered good for digestion and overall health.

Others have implicated probiotics in the production of D-lactic acid -- and brain foginess -- in patients with a short bowel so their small intestine does not function properly, and in newborns fed formula containing the popular product. Short bowel syndrome results in a lot of undigested carbohydrates that are known to cause small intestinal bacterial overgrowth, or SIBO, and the high levels of D-lactic acid. Severe liver and kidney problems can produce similar problems.

Whether there was also a connection when the gut is intact was an unknown. "This is the first inroad," says Rao.

All patients experiencing brain foginess took probiotics and SIBO was more common in the brain foginess group as well, 68 percent compared to 28 percent, respectively. Patients with brain foginess also had a higher prevalence of D-lactic acidosis, 77 versus 25 percent, respectively.

When brain-foggy patients stopped taking probiotics and took a course of antibiotics, their brain foginess resolved.

Movement of food through the gastrointestinal tract was slow in one third of the brain foggy patients and one fourth of the other group. Slower passage, as well as things like obesity surgery, can increase the chance of bacterial buildup, or SIBO.

"Now that we can identify the problem, we can treat it," Rao says. Diagnosis includes breath, urine and blood tests to detect lactic acid, and an endoscopy that enables examination of fluid from the small intestines so the specific bacteria can be determined and the best antibiotics selected for treatment.

Normally there is not much D-lactic acid made in the small intestines, but probiotic use appears to change that. SIBO, which was present in most with brain foginess, can cause bacteria to go into a feeding frenzy that ferments sugars resulting in production of uncomfortable things like hydrogen gas and methane that explain the bloating.

Probiotics added to that feeding frenzy the bacterium lactobacillus, which produces D-lactic acid as it breaks down sugars, The acid get absorbed in

Electroacupuncture Reduces Inflammation Via Specific Neural Pathways

Nanotubes in the Eye That Help Us See

MIND & BRAIN

Evolutionary Theory of Economic Decisions



Neanderthals May Have Had a Lower Threshold for Pain

Antibiotics Disrupt Development of the 'Social Brain' in Mice

LIVING & WELL

'Drawn-on-Skin' Electronics Offer Breakthrough in Wearable Monitors

New Fabric Could Help Keep You Cool in the Summer, Even Without A/C

Giving Robots Human-Like Perception of Their Physical Environments

the blood and can reach the brain.

All those with brain fogginess, SIBO and/or D-lactic acidosis, were given antibiotics that targeted their bacterial population and asked to discontinue probiotics. Those without SIBO were asked to halt probiotics and stop eating yogurt, which is considered one of the best sources of probiotics. Those with SIBO and D-lactic acidosis but no brain fogginess also took antibiotics.

Following treatment, 70 percent of patients reported significant improvement in their symptoms and 85 percent said their brain fogginess was gone.

Those without brain fogginess but with SIBO and high levels of D-lactic acid reported significant improvement in symptoms like bloating and cramping within three months.

Abdominal pain was the most common symptom in both groups and before treatment, six of those with brain fogginess reported a tremendous increase in their abdominal size within just a few minutes of eating.

All patients received extensive examination of their gastrointestinal tract, including a motility test, to rule out other potential causes of their symptoms. They filled out questionnaires about symptoms like abdominal pain, belching and gas and answered questions about related issues like antibiotic and probiotic use as well as food fads and yogurt consumption.

They were given carbohydrates followed by extensive metabolic testing looking at the impact on things like blood glucose and insulin levels. Levels of D-lactic acid and L-lactate acid, which results from our muscles' use of glucose as energy and can cause muscle cramps, also were measured.

Probiotic use may be particularly problematic for patients who have known problems with motility, as well as those taking opioids and proton pump inhibitors, which reduce stomach acid secretion and so the natural destruction of excessive bacteria.

Probiotics are supposed to work in the colon and not the small intestines or stomach, Rao says, so motility issues can result in problems with probiotic bacteria reaching the proper place. A wide variety of problems, from conditions like diabetes to drugs like antidepressants and minerals like iron, can slow movement and increase the possibility that probiotics will remain too long in the upper gut where they can cause harm, he says.

Probiotics definitely can help, for example, people who have gastroenteritis, or stomach flu, or are left with diarrhea and other problems after antibiotics wipe out their natural gut bacteria, Rao says.

"In those situations, we want to build up their bacterial flora so probiotics are ideal," he says.

Rao's pursuit of a possible connection between probiotics, brain fogginess and bloating started with a memorable patient who developed significant amounts of both problems within a minute of eating.

"It happened right in front of our eyes," Rao says of the dramatic abdominal distention. They knew the woman had diabetes, which can slow motility. When they looked in the blood and urine at a variety of metabolic com-

pounds, they found the high levels of D-lactic acid and soon learned the patient used probiotics and regularly ate yogurt.

Next steps include additional studies in which the investigators better quantify and characterize the brain fogginess reported by patients and following patients for longer periods to ensure their problems remain resolved. Some patients in the current study required a couple of rounds of antibiotics, Rao notes.

Good food sources of probiotics include yogurt, sauerkraut, kimchi, kefir and dark chocolate, which are generally safe because of the small amounts of bacteria present, Rao says.

The 19-foot long small intestine has been a bit of an understudied organ, likely in part because it's hard to visualize via the mouth or anus, Rao says. "I think the small bowel can be a source of huge mystery," Rao says.

Your helpful gut bacteria, or microbiome, which are essential to things like a well-functioning immune system and general health, are largely in the large intestine and colon.

Story Source:

Materials provided by **Medical College of Georgia at Augusta University**.
Note: Content may be edited for style and length.

Journal Reference:

1. Satish S. C. Rao, Abdul Rehman, Siegfried Yu, Nicole Martinez de Andino. **Brain fogginess, gas and bloating: a link between SIBO, probiotics and metabolic acidosis.** *Clinical and Translational Gastroenterology*, 2018; 9 (6) DOI: 10.1038/s41424-018-0030-7

Cite This Page:

MLA	APA	Chicago
-----	-----	---------

Medical College of Georgia at Augusta University. "Probiotic use is a link between brain fogginess, severe bloating." ScienceDaily. ScienceDaily, 6 August 2018. <www.sciencedaily.com/releases/2018/08/180806095213.htm>.

RELATED STORIES

Protecting Probiotics from the Stomach

Sep. 26, 2018 — The small intestine is a hotbed of microbial activity and a target of probiotic treatments for diarrhea, inflammatory bowel disease and irritable bowel syndrome, among other conditions. To make it to ...

Lack of 'Editing' in Brain Molecules Potential Driver of Cancer

June 16, 2017 — A “significant” lack of ‘editing’ has been observed in microRNAs in brain tissue of brain cancer patients, report ...

A Unique Enzyme Could Be a Game-Changer for Gluten-Sensitive Patients
May 8, 2017 — Researchers have found that taking an enzyme tablet while consuming foods containing gluten prevents a significant amount of it from entering the small intestine. This could enable gluten-sensitive ...

Taking a Combination of Painkillers, Gastric Protection Damages the Intestines

Apr. 11, 2017 — Patients with inflammatory diseases are often prescribed non-steroidal anti-inflammatory drugs. They are also often recommended to use a proton pump inhibitor to protect their stomach. In a new ...

FROM AROUND THE WEB

Below are relevant articles that may interest you. ScienceDaily shares links with scholarly publications in the TrendMD network and earns revenue from third-party advertisers, where indicated.

Work Together

GenomeWeb, 2011

Utilizing Evolutionary Mechanisms of Human-Bacterial Symbiosis

Editors, Medgadget, 2008

Depression in oncologists: For many, a closely guarded secret

Anthony L. Back et al., Healio, 2019

The Way to a Mouse's Brain Is Through its Stomach

GenomeWeb, 2011

Catalog of Mobile Antibiotic Resistance Genes Catalogued in Bacteria Helps Explain Transfer to Human Pathogens

GenomeWeb, 2016

Importance of gut microbiota in obesity

Medicine Matters diabetes, 2018

New Biomarker of Intestinal Permeability Could Lead to Dx for Gut Function

Christie Rizk, GenomeWeb, 2019

Clinical value of MRD detection in blood cancers ‘increasingly difficult to ignore’

Timothy S. Pardee et al., Healio, 2013

Powered by **TREND MD**

Free Subscriptions

Get the latest science news with ScienceDaily's free email newsletters, updated daily and weekly. Or view hourly updated newsfeeds in your RSS reader:

 [Email Newsletters](#)

 [RSS Feeds](#)

Follow Us

Keep up to date with the latest news from ScienceDaily via social networks:

 [Facebook](#)

 [Twitter](#)

 [LinkedIn](#)

Have Feedback?

Tell us what you think of ScienceDaily -- we welcome both positive and negative comments. Have any problems using the site? Questions?

 [Leave Feedback](#)

 [Contact Us](#)

[About This Site](#) | [Staff](#) | [Reviews](#) | [Contribute](#) | [Advertise](#) | [Privacy Policy](#) | [Editorial Policy](#) | [Terms of Use](#)

Copyright 2020 ScienceDaily or by other parties, where indicated. All rights controlled by their respective owners. Content on this website is for information only. It is not intended to provide medical or other professional advice. Views expressed here do not necessarily reflect those of ScienceDaily, its staff, its contributors, or its partners.

Financial support for ScienceDaily comes from advertisements and referral programs, where indicated.

— [CCPA: Do Not Sell My Information](#) — — [GDPR: Privacy Settings](#) —